Women's and Children's Health



Current General MCH Priorities

- Teen pregnancy/family planning
- Infant mortality
- Obesity/overweight
- Injury prevention
- Oral health
- Prenatal care
- Mental health



Health Indicators

Indicators for three populations:

- 1) Children and Adolescents
- 2) Pregnant women and infants
- 3) Women of Reproductive Age (15-44 Years)



Child & Adolescent Health State	us, Well-Being and F	Protectiv	e Fact	tors

Indicators include:	Estimate	Comparison / AZ Trend
2-4 years with untreated tooth decay (2009)	31.0%	16.0% (U.S.)
3 rd grade students with untreated tooth decay (2009)	41.0%	29.0% (U.S.)
0-17 year old without health insurance (2008)	16.0%	9.9% (U.S.) (2003-2008)
19-35 month old children with 4:3:1:3:3 vaccines (2007)	79.2%	78.2% (U.S.) (2004-2008)
2 to <5 year old low income children who are obese (2008)	14.6%	14.9% (U.S.) (2003-2008)
0-17 year old preventable deaths (2008)	33.0%	(2006-2008)

Child & Adolescent Health Status, Well-Being and Protective Factors			
Indicators include:	Estimate	Comparison / AZ Trend	
15-19 year old teen pregnancy (2008)	63.2 per 1,000	(1999-2008)	
15-19 year old teen births (2008)	54.9 per 1,000	42.5 (U.S.) (1999-2008)	
15-17 year old teen births w/1st trimester PNC (2008)	63.5%	90.0% (U.S.) (2003-2008)	
18-19 year old teen births w/1st trimester PNC (2008)	69.7%	90.0% (U.S.) (2003-2008)	
15-19 year old teen births w/ one or more medical risk factors (2008)	34.6%	(2003-2008)	

25.9%

16.1%

22.2%

28.8% (U.S.)

14.5% (U.S.)

20.0% (U.S.)

(2003-2007)

High school students who are overweight or

High school students who seriously

considered attempting suicide (2007)

High school students who currently smoke

obese (2007)

tobacco (2007)

Pregnant Women and Infant Health Status, Well-Being and Protective Factors

Indicators include:	Estimate	Comparison / AZ Trend
1 st trimester prenatal care (2008)	79.4%	84.0% (U.S.) (2003-2008)
Preterm births (<37 weeks gestation) using clinical estimate of gestation (2008)	10.2%	7.6% (HP 2010) (1999-2008)
Low birth weight (<2,500 grams) infant (2008)	7.1%	8.3% (U.S.) (1999-2008)
Stillbirth rate (>20 weeks gestation) (2008)	5.5 per, 1,000	6.2 (U.S.) (2003-2008)
Perinatal mortality rate (28 weeks gestation to < 7 days age) (2008)	6.3 per 1,000	6.6 (U.S.) (1999-2008)
Infant mortality rate (2008)	6.3 per 1,000	6.5 (U.S.) (1999-2008)

Pregnant Women and Infant Health Status, Well-Being and Protective Factors

Indicators include:	Estimate	Comparison / AZ Trend
SIDS rate (2008)	4.0 per 10,000	5.1 (U.S.) (1999-2008)
Women who reported smoking during pregnancy (2008)	4.8%	1.0% (HP 2010) (1999-2008)
Women age 20-34 years with one or more medical risk factors (2008)	30.3%	(2003-2008)
Primary cesarean section for singleton deliveries at term (37-41 weeks) (2008)	21.7%	15.0% (HP 2010) (1999-2008)
Macrosomic infant (>4,000 grams) (2008)	7.3%	(2003-2008)
Maternal mortality rate (< =42 days postpartum) (2008)	10.1 per 100,000	3.3 (HP 2010) (2000-2008)
Any breastfeeding at 6 months postpartum (2008)	45.3%	43.4% (U.S.) (2005-2008)

Women of Reproductive Age Health Status, Well-Being, Protective Factors

Indicators include:	Estimate	Comparison / AZ Trend
Obesity women 18-44 years (2008)	22.3%	15.0% (HP 2010) (2000-2008)
Take folic acid supplement or vitamin women 18-44 years (2007)	50.2%	80.0% (HP 2010) (2004-2007)
Eat "5-a-day" vegetables and fruit women 18-44 years (2008)	25.5%	50.0% (HP 2010) (2003-2008)
Get recommended physical activity per week women 18-44 years (2008)	58.1%	50.0% (HP 2010) (2003-2008)
Report frequent mental distress women 18-44 years (2008)	19.8%	12.0% (U.S.) (2003-2008)
Chlamydia incidence women 15-24 years (2008)	3010 per 100,000	1650 (HP 2010) (1999-2008)
Gonorrhea incidence women 15-24 years (2008)	235.1 per 100,000	270 (HP 2010) (1999-2008)
Forcible rape incidence (2008)	25.7 per 100,000	29.3 (U.S.) (2003-2008)


Children and Adolescents Health Status and Well-Being Indicators

Examples:

- Mortality and injury
- Mental health
- Oral health
- Chronic Disease




Mortality Rates for Leading Causes of Death per 100,000 Children Age 1-14 Years, Arizona 1999-2008



*significant difference between 1999 and 2008 at α =0.05 Source: Arizona Vital Statistics, 1999-2008



Mortality Rate for All Deaths per 100,000 Children Age 1-14 Years by Area of Residence, Arizona 1999-2008




*significant difference between the urban and rural rates at α=0.05
Urban areas are; Maricopa, Pima, Pinal, and Yuma Counties. All other counties are considered rural.

Source: Arizona Vital Statistics. 1999-2008




Mortality Rates for Leading Causes of Death per 100,000 Adolescents Age 15-19 Years, Arizona 1999-2008



^{*}significant difference between 1999 and 2008 rates at α =0.05 Source: Arizona Vital Statistics, 1999-2007




Mortality Rates for All Causes of Death per 100,000 Adolescents Age 15-19 Years by Area of Residence, Arizona 1999-2008



^{*} significant difference in mortality rates between urban and rural residents at α =0.05 Source: Arizona Vital Statistics, 1999-2008



Female High School Students Reporting Seriously Considered Attempting Suicide, Arizona 2003, 2005, 2007




Source: Youth Risk Behavior Survey, 2003, 2005, 2007






Male High School Students Reporting Seriously Considered Attempting Suicide Arizona & U.S. 2003, 2005, 2008



*significant difference between AZ and U.S. at α =0.05 Source: Youth Risk Behavior Survey, 2003, 2005, 2007




Mortality Rate for Deaths by Suicide per 100,000 Adolescents Age 15-19 Years Arizona & U.S. 2003-2008



^{*}significant difference between AZ and the U.S. at α =0.05



Children Age 2-4 Years with Untreated Tooth Decay U.S. & Arizona, 2009




Source: Arizona Healthy Smiles Health Bodies Survey, 2009 (data not weighted to population)





3rd Grade with Untreated Tooth Decay U.S. & Arizona, 2009




Source: Healthy Smiles Healthy Bodies Survey, 2009 (data not weighted to population)


Leadership for a Healthy Arizona



Hospitalization Rate for Asthma per 10,000 Children Age <5 Years by Race/Ethnicity, Arizona 2004-2008




Prevalence of Current Asthma Diagnosis Among High School Aged Students Arizona & U.S. 2007



Source: Arizona YRBS, 2007



Diabetes-related Emergency Room Visits and Inpatient Discharges per 10,000 Children and Adolescents, Arizona 2005-2008



Source: Arizona Hospital Discharge Data, 2005-2008




Children and Adolescents Health Determinants, Risk, and Protective Factors

Examples:

- Overweight and obesity
- Teen pregnancy and birth
- Substance abuse/Anti-social behavior



Prevalence of Obesity and Overweight among Low Income Children Age 2 to <5 Years, Arizona 2004-2008



Source: Pediatric Nutrition Surveillance, 2008





TABLE 1
Prevalence of Overweight and Obesity by Race/Ethnicity of Low Income Children Age 2 to <5 Years
Arizona, 1999 & 2008

White non-Hispanic	1999¹	20081	Percent Change
Overweight	10.9	14.0	28.4
Obesity	6.6	9.5	43.9
Total Overweight or Obese	17.5	23.5	34.3
Hispanic or Latino			
Overweight	13.2	16.6	25.8
Obesity	11.5	16.2	40.9
Total Overweight or Obese	24.7	32.8	32.8
Black or African American			
Overweight	10.4	13.5	29.8
Obesity	6.5	10.3	58.5
Total Overweight or Obese	16.9	23.8	40.8
American Indian or Native			
Overweight	15.2	20.4	34.2
Obesity	15.7	20.1	28.0
Total Overweight or Obese	30.9	40.5	31.1
Asian or Pacific Islander			
Overweight	9.9	12.1	21.8
Obesity	8.5	14.4	69.4
Total Overweight or Obese	18.4	26.5	44.0

Source: Pediatric Nutrition Surveillance, 2008


1 per 100 children age 2 to <5 years.

Leadership for a Healthy Arizona

Arizona

Department of
Health Services

Percent of Obesity and Overweight among Low Income Children Age 2 to <5 Years Arizona Counties, 2006-2008




Source: Pediatric Nutrition Surveillance, 2008





Prevalence of Obesity or Overweight Among High School Aged Youth Arizona, 2003/2005/2007




■ White non-Hispanic □ Hispanic or Latino ■ American Indian or Alaska Native

Source: Youth Risk Behavior Survey 2003, 2005, 2007





Percent of Children Age 10-17 at 'Healthy Weight' by Family Income Level Arizona 2003, 2007




FPL= Federal Poverty Level

*significantly greater percentage of children at healthy weight compared to children at 100%-199% FPL at α=0.05 Source: National Survey of Children's Health, 2003, 2007

Leadership for a Healthy Arizona



Estimated Teen Pregnancy Rates Females Age 15-19 Years Arizona 1999-2008




Source: Arizona Vital Statistics, 1999-2008




Leadership for a Healthy Arizona

Teen Pregnancy for Females Age 15-19 Years by Race/Ethnicity, Arizona 1999-2008



Health Services


Teen Pregnancy for Females Age 15-19 Years by Urban and Rural Residence, Arizona 2003-2008







Births to Females Age 15-19 Years Arizona and the U.S. 1999-2008




Source: Arizona Vital Statistics, 1999-2008




Leadership for a Healthy Arizona


Birth Rates per 1,000 Females Age 15-17 Years by Arizona Community Health Analysis Area (CHAA), 2008 Kaibab Paiute Fredonia Navajo Nation Littlefield Williams Flagstaff-Chino Valley/ ad City Bullh Kingman Ash Fork Fort I Holbrook St. Johns ke Hayasu Heber-Overgaai Snowflake Yavapai Co. S/Bagdad Corde Payson Colorado Indian Wickenburg Graham Co. S Yuma NW Willcox/Bowie Arivaca Continent Birth Rates by Quartile Tombstone/Elfrida 1.8 to 17.4 county boundary 17.5 to 30.5 30.6 to 49.2 Arizona Department of 49.3 and above Health Services Source Data: Arizona Birth Certificates and Arizona Office of Vital Statistics, 2008

Metro Phoenix




Metro Tucson






Percent of Births to Females Age 15-19 Years by Time of Prenatal Care Initiation, Arizona 2003-2008



Source: Arizona Vital Statistics, 2003-2008




Mothers with Medical Risk Factors per 100 Live Births for Females Age 15-19 and Age 20-34 Years, Arizona 2003-2008



* significant difference at α=0.05 Source: Arizona Vital Statistics, 2003-2008




Alcohol Consumption in Past Month by High School Aged Students Arizona & U.S. 2003/2005/2007



Leadership for a Healthy Arizona




Smoked Cigarettes in Past Month by High School Aged Students Arizona & U.S. 2003/2005/2007






Lifetime Use of Methamphetamine by High School Aged Students Arizona & U.S. 2003/2005/2007






Lifetime Substance Use Among 8th, 10th, & 12th Grade Students Arizona Youth Survey, 2008







Delinquency and Anti-Social Behavior Arizona Youth Survey, 2008





Pregnant Women and Infants




Pregnant Women and Infants Health Status and Well-Being Indicators

Examples include;


- 1) Preterm birth and low birth weight
- 2) Infant mortality/Stillbirths








Percent Change in Total Resident Births from 2007 to 2008 Arizona 2008








Percent of Total Births by Mother's Race/Ethnicity Arizona 2008



Source: Arizona Vital Statistics, 2008



Percent of Infants Born Preterm (<37 weeks gestation) by Type of Gestational Estimate Arizona 1999-2008



^{*}prematurity based on physicians estimate of gestational age

Source: National Center for Health Statistics and Arizona Vital Statistics, 2008




^{**}prematurity based on mother's recall date of last menses

Preterm Births (< 37 weeks)*, Arizona 2003-2008										
Race/Ethnicity	2003	2004	2005	2006	2007	2008				
White non-Hispanic	11.0%	11.4%	10.9%	10.5%	10.4%	10.1%				
Black/African American	11.7%	11.6%	12.6%†	12.7%†	12.8%†	11.7%†				
American Indian/Alaskan Native	9.8%†	10.9%	10.6%	11.7%†	10.4%	9.8%				
Asian/Pacific Islander	10.8%	11.0%	10.2%	10.4%	9.6%	10.0%				
Hispanic/Latina	9.8%†	10.3%†	10.1%†	10.1%	9.8%	9.9%				
Mother's Age										
18-34 years	10.0%	10.5%	10.2%	10.1%	9.9%	9.7%				
<18 years	10.9%	11.8%†	12.1%†	11.9%†	10.7%	11.0%†				
40 years +	16.9%†	17.6%	16.8% [†]	15.0% [†]	16.4% [†]	17.2%†				
Number of Births										
Singleton	8.9%	9.2%	9.2%	9.0%	8.7%	8.5%				
Multiple	64.7%†	70.3%†	64.6% [†]	67.6% [†]	67.4% [†]	65.5% [†]				
Payee for Birth										
Private Insurance	10.5%	10.8%	10.5%	10.3%	10.2%	10.1%				
AHCCCS	10.5%	11.4%†	10.9%	10.8%	10.4%	10.2%				
Area of Residence										
Urban County	10.6%	11.0%	10.8%	10.7%	10.3%	10.2%				
Rural County	10.0%	10.9%	10.0%†	10.0%†	10.0%	9.7%				

^{*}Physician's estimate of gestational age

Italicized category is reference group † statistically significant difference from reference group at α <0.05 Source: Health Status and Vital Statistics, 2003-2008

Low Birth Weight (<2,500 grams) and Very Low Birth Weight (<1,500 grams) U.S. and Arizona 1999-2008



^{*}latest year of national data available at time of data collection.



Low Birth Weight (< 2,500 grams), Arizona 2003-2008										
Race/Ethnicity	2003	2004	2005	2006	2007	2008				
White non-Hispanic	6.9%	7.2%	6.9%	6.7%	7.1%	6.9%				
Black/African American	11.7%†	11.6%†	12.6%†	12.7%†	12.8%†	11.7%†				
American Indian/AK Native	6.8%	7.2%	7.3%	7.4%	6.6%	6.7%				
Asian/Pacific Islander	8.5%†	8.0%	8.0%	8.5%†	8.0%	9.1%†				
Hispanic/Latina	6.9%	6.8%	6.4%	6.9%	6.6%	6.7%				
Mother's Age										
18-34 years	6.8%	6.9%	6.7%	6.8%	6.8%	6.7%				
<18 years	8.4%†	8.4%†	8.4%†	8.5%†	8.3%†	9.1%†				
40 years +	12.4%†	12.2%†	9.6%†	9.6%†	11.4%†	11.4%†				
Number of Births										
Singleton	5.6%	5.6%	5.5%	5.7%	5.6%	5.6%				
Multiple	58.3% [†]	59.7% [†]	56.3% [†]	57.8% [†]	60.3% [†]	57.0%†				
Payee for Birth										
Private Insurance	6.6%	6.7%	6.5%	6.6%	6.8%	6.9%				
AHCCCS	7.4%†	7.5%†	7.2%†	7.5%†	7.4%†	7.2%				
Prenatal Care										
1 st Trimester	6.9%	7.1%	6.9%	7.1%	7.1%	7.1%				
Other or None	8.5%†	7.4%	8.4%†	9.1%†	9.1%†	8.9%†				
Area of Residence										
Urban County	7.0%	7.1%	6.9%	7.0%	7.0%	7.0%				
Rural County	7.3%	7.7%	7.4%	7.7%	7.5%	7.5%				


Italicized category is reference group

Source: Health Status and Vital Statistics, 2003-2008

 $[\]ensuremath{^{**}}$ Urban Counties are Maricopa, Pinal, Pima, and . All other counties are considered rural.

 $^{^{\}text{+}}\text{statistically significant difference}$ from reference group at α <0.05

Rate of Infant Deaths per 1,000 Live Births Arizona and the United States, 1999-2008




*Provisional rates for the U.S. in 2007 and 2008 Source: Arizona Health Status and Vital Statistics, 2008






Rate of Infant Deaths per 1,000 Live Births by Race and Ethnicity Arizona, 1999-2008






Proportion of Total Infant Deaths by Race and Ethnicity Arizona, 1999-2008 (n= 6,083)







Infant Mortality Rates for the Five Leading Causes of Death by Race and Ethnicity, Arizona, 2008



- ■Sudden infant death syndrome
- Complications of placenta, cord and membrane
- Maternal complications
- ■Short gestation and low birth weight
- Congenital malformations, deformations and chromosomal abnormalities





Neonatal and Postneonatal Mortality Rates and Percent Postneonatal Deaths 1999-2008 **UNITED STATES** 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 4.7 4.5 Neonatal mortality 4.7 4.6 4.6 4.6 4.6 4.5 NA NA 2.3 2.3 2.3 2.3 2.3 Postneonatal mortality 2.3 NA NA 33.0 33.3 33.8 33.3 33.3 Percent postneonatal mortality 33.3 32.8 32.8 NA NA ARIZONA STATEWIDE Neonatal mortality 4.5 4.3 4.5 4.1 4.3 4.5 4.4 4.3 4.7 4.2 2.2 2.3 Postneonatal mortality 2.4 2.4 2.2 2.4 2.0 2.2 2.1 33.8 35.9 35.0 33.8 32.8 35.3 31.5 32.4 33.3 Percent postneonatal mortality 34.8 **URBAN COUNTIES** 4.7 Neonatal mortality 4.5 4.2 4.7 4.2 4.4 4.3 4.3 4.6 4.3 2.2 2.3 2.3 Postneonatal mortality 2.2 2.0 2.0 2.2 2.0 2.0 2.0 33.8 33.8 Percent postneonatal mortality 35.4 31.9 34.4 31.5 29.9 31.3 30.3 31.8 **RURAL COUNTIES** 4.6 3.4 3.8 3.8 3.8 5.0 4.3 4.8 3.5 Neonatal mortality 4.6 2.1 3.6 3.3 2.7 Postneonatal mortality 3.1 2.9 2.4 3.3 3.1 2.0 38.7 43.6 31.3 40.3 46.0 46.2 44.9 41.9 32.6 41.3 Percent postneonatal mortality

Source: Health Status and Vital Statistics, 2008


Note: Urban = Maricopa, Pima, Pinal and . The remaining counties comprise 's rural areas. Records with unknown county of residence are included in the statewide totals, but are not distributed by urban/rural area.

Arizona

Department of Health Services



Percent of Infant Mortality Rate that is Postneonatal Mortality Arizona's Urban and Rural Counties, 1999-2008



Year

■ URBAN COUNTIES ■ RURAL COUNTIES

Source: Arizona Health Status and Vital Statistics, 2008


Urban Counties are Maricopa, Pima, Pinal and Yuma. The remaining counties comprise Arizona's rural areas.





^{*} Significantly different at p<0.05.


Stillbirth Rates (per 1,000 live births + fetal deaths*) Arizona 2003-2008 (n= 3,246)



*fetal deaths of 20 weeks or more gestation Source: Arizona Fetal Death Certificate Data, 2003-2008




Stillbirth Rates (per 1,000 live births+fetal deaths) by Race/Ethnicity, Arizona 2003-2008 (n= 3,231)*



*excludes unknown race/ethnicity Source: Arizona Fetal Death Certificate Data. 2003-2008










Pregnant Women and Infants Health Determinants, Risk and Protective Factors

Examples include;

- 1) Tobacco use during pregnancy
- 2) Cesarean Deliveries
- 3) Medical conditions during pregnancy
- 4) Breastfeeding




Reported Smoking During Pregnancy by Race/Ethnicity Arizona 2008



Source: Arizona Vital Statistics, 2008




Mothers Who Reported Smoking Tobacco During Pregnancy by Insurance Status Arizona 1999-2008



*significant difference at α =0.05 Source: Arizona Birth Certificates, 1999-2008




Primary Cesarean Sections for all Term (37-41 weeks)* and Preterm Infants (<37 weeks)*, Arizona, 1999-2008





Primary Cesarean Sections for Singleton Term Births (37-41 weeks) by Race/Ethnicity, Arizona, 1999-2008




Source: Arizona Birth Certificate data, 1999-2008





Primary Cesarean Sections for Singleton Term Births (37-41 weeks) by Insurance Status. Arizona 1999-2008




*significant difference between private insurance and public insurance at α =0.05 Source: Arizona Birth Certificate data, 1999-2008





Vaginal Birth After Cesarean Section Arizona 1999-2008




Source: Arizona Birth Certificate Data, 1999-2008






Any Diabetes During Pregnancy per 1,000 Live Births Arizona 2003-2008



*significant difference at α=0.05 Source: Arizona Vital Statistics, 2003-2008




Anemia During Pregnancy per 1,000 Live Births Arizona 2003-2008






Reported Congenital Syphlis Cases and Case Rates per 100,000 Live Births by Birth Year,





Breastfeeding at 6-Months for Infants in Low Income Families Arizona (2008) & U.S. (2007)




Source: Pediatric Nutrition Surveillance System, 2008





Breastfeeding at 6-Months for Infants in Low Income Families Arizona, 2004-2008



Source: Pediatric Nutrition Surveillance System, 2008





Women of Reproductive Age




Women of Reproductive Age Health Status and Well-Being Indicators

Examples include;

- 1) Mortality
- 2) Mental health
- 3) STDs




Mortality Rates for the Leading Causes of Death Among Women Age 20-44 Years, Arizona 1999-2008



Health Services

Leadership for a Healthy Arizona


Mortality Rate for Leading Causes of Death Among Women Age 20-44 Years by Urban and Rural Residence, Arizona 1999-2008



Source: Arizona Vital Statistics, 1999-2008




Frequent Mental Distress for Women Age 18 thru 44 Years by Race/Ethnicity, Arizona 2003-2008



Source: Arizona BRFSS, 2003-2008



Obesity Prevalence Among Females Age 18-44 Years With and Without Frequent Mental Distress, Arizona 2003-2008




*significant difference at α=0.05 Source: Arizona BRFSS, 2003-2008

Leadership for a Healthy Arizona




Incidence of Chlamydia among Females by Age Group Arizona (2008) & U.S. (2007)



Source: Arizona Vital Statistics and the CDC, 2008



Reported Chlamydia Case Rates per 100,000 Population by Gender Arizona 2003-2008




Leadership for a Healthy Arizona

Arizona

Department of
Health Services

Reported Gonorrhea Case Rates per 100,000 Population by Gender Arizona 2003-2008




Leadership for a Healthy Arizona

Arizona

Department of
Health Services


Reported Primary and Secondary Syphilis Case Rates per 100,000 Population by Gender, Arizona 2003-2008



Leadership for a Healthy Arizona

Arizona
Department of
Health Services

HIV/AIDS Rate By Age at Emergent Infection Among Women by Race/Ethnicity Arizona, 2003-2007




Women of Reproductive Age Health Determinants, Risk and Protective Factors

Examples include;

- 1) General Health/Obesity
- 2) Folic Acid
- 3) Smoking
- 4) Violence




Percent of Females Age 18-44 Years Self-Reported Health Status by Household Income Level, Arizona 2001-2008



^{*}significant difference at α =0.05. Percentages weighted to population Source: Arizona BRFSS, 2001-2008



Percent of Women Aged 18-44 Reporting No Leisure Time Physical Activity in Past 30 Days, Arizona 2008




Source: Arizona BRFSS, 2008 (percentages weighted to population)



Leadership for a Healthy Arizona

Percent of Women Age 18-44 Years Eating "5-A-Day" by Race/Ethnicity Arizona 2003-2008




Source: Arizona BRFSS, 2003-2008 (percentages weighted to population)





Prevalence of Obesity Women Aged 18 thru 44 Arizona 2000-2008




*significant difference between point estimate and HP 2010 Goal Source: Arizona BRFSS, 2000-2008 (percentages weighted to population)






Percent of Females Aged 18 thru 44 who are Obese by Urban & Rural Regions, Arizona 2000-2008



*significant difference between rural obese estimate and HP 2010 Goal Source: Arizona BRFSS, 2000-2008



Percent of Women Age 18-44 Years Consuming a Vitamin or Supplement with Folic Acid by Level of Education, Arizona 2005-2007




^{*}significant difference (p<0.05)

Source: Arizona BRFSS 2005-2007 (percentages weighted to population)






Current Smokers among Women Age 18-44 Years by Race/Ethnicity Arizona 2003-2008








Rate of Homicide for Female Victims Age 20-44 Years Arizona 2003-2008





Rate of Forcible Rape Against Females of All Ages Arizona & U.S. 2004-2008



Source: FBI UCR, 2004-2008





Adequacy of Services Provided in Your Community for Selected MCH Populations

Children and Teenagers	Very well or sometimes well	Not well or never
Safe places to play and exercise (n=764)	54.8%	45.2%
Reproductive health/family planning (n=717)	54.7%	45.3%
Mental health care (n=732)	45.0%	55.1%
Drug and alcohol treatment (n=720)	43.8%	56.3%
Teen driver and safety training (n=665)	39.9%	60.2%
After school activities for teens (n=719)	39.9%	60.2%



Adequacy of Services Provided in Your Community for Selected MCH Populations

Pregnant Women and Infants	Very well or sometimes well	Not well or never
Access to hospital with delivery facilities (n=757)	79.9%	20.1%
Birth education classes (n=719)	75.2%	24.8%
Breastfeeding education (n=746)	72.0%	28.0%
Dental care (n=738)	37.7%	62.3%
Affordable transportation to access health care (n=711)	37.7%	62.3%
Support for breastfeeding at work (n=712)	31.7%	68.3%



Adequacy of Services Provided in Your Community for Selected MCH Populations

Women of Reproductive Age (15-44 years)	Very well or sometimes well	Not well or never
Reproductive health and family planning (n=754)	72.3%	27.7%
Affordable and healthy food for people using food stamps (n=741)	67.3%	32.7%
Shelters for domestic violence victims (n=742)	65.2%	34.7%
Mental health treatment (n=727)	47.5%	52.5%
Affordable health care (n=755)	43.6%	56.4%
Dental care (n=735)	42.5%	57.6%

